

DUANE M. GELS, MD, FAAAAI Board Certified ABAI Clinical Faculty – Johns Hopkins 129 Lubrano Drive, Suite 200 Annapolis, MD 21401 Phone 410.573.1600 Fax 410.573.5841 www.annapolisallersy.com

# PREPARATION FOR ALLERGY TESTING

## MEDICATIONS YOU ARE REQUIRED TO DISCONTINUE:

ANTIHISTAMINES interfere with skin testing. Discontinue all prescription and over-the-counter antihistamines at least 7 days prior to testing. Below is a list of common antihistamines. Check your medication label for these. This list may not be complete. If you are unsure, please CALL OUR OFFICE AT 410.573.1600.

Advil Allergy Advil PM Alavert Alka-Seltzer Plus Allegra Allegra D Allegra ODT Astelin Astepro Atarax Benadryl Bromfed Brompheniramine Cetirizine Chlorpeniramine Chlortrimeton Clarinex Clarinex Clarinex D Claritin Claritin Claritin D Comtrex Diphenhydramine

Desloratadine Dimetapp Doxylamine Dymista Fexofenadine Hydroxyzine Levocetirizine Loratadine Nyquil "PM" label (ANY BRAND) Patanase Phenergan Promethazine Robitussin Theraflu Triaminic Tussionex Tylenol PM Xyzal Zyrtec Zyrtec D

### MEDICATIONS YOU MAY OF MAY NOT BE REQUIRED TO DISCONTINUE:

BLOOD PRESSURE, HEART, DEPRESSION, ORAL STEROIDS (PREDNISONE) or MIGRAINE MEDICATIONS <u>may</u> affect testing. <u>Do not discontinue these medications without consulting our office</u>.

### DO NOT DISCONTINUE ASTHMA MEDICATIONS:

Examples: Singulair, Advair, Alvesco, Asmanex, Dulera, Symbicort, Flovent, Pulmicort, QVAR, Albuterol (Ventolin, Proventil, ProAir, Xopenex), and any other lung inhalers.

# IF YOU ARE UNSURE ABOUT ANY MEDICATION, PLEASE CALL 410.573.1600.



DUANE M. GELS, MD, FAAAAI Board Certified ABAI Clinical Faculty – Johns Hopkins 129 Lubrano Drive, Suite 200 Annapolis, MD 21401 Phone 410.573.1600 Fax 410.573.5841 www.aunapolisallerpy.com

# PREPARATION FOR ALLERGY TESTING

### ADDITIONAL INSTRUCTIONS:

- DO NOT APPLY CREAMS OR LOTIONS TO ARMS unless medically necessary. Deodorant/Antiperspirant is OK.
- Please WEAR or BRING A LOOSE, SHORT SLEEVED SHIRT on the day of testing. We will need access to your arms up to your shoulders.
- LENGTH OF VISIT cannot always be predicted. Be prepared to stay 2 4 hours, unless we advise you otherwise. Feel free to bring a book or a DVD. A DVD player is available.
- EATING prior to visit is recommended as food is not allowed in the office. (see below)
- BRING COPIES of Blood Tests (if relevant), previous ALLERGY TESTS, Chest X-RAYS, and/or CT Scans.

#### IN CASE OF ACUTE ILLNESS:

IF YOU ARE ACTIVELY WHEEZING, COUGHING, or incidentally have HIVES (welts) on the day of your visit, allergy testing may not be possible. If this is your first visit, will be evaluated by Dr. Gels, but testing may need to be postponed. Alternatively, you may call us to reschedule.

# II VERY IMPORTANT II PLEASE I NO FOOD OR DRINK IN OFFICE I

Young patients with life-threatening food allergies are regular visitors in our office. A small crumb found by a toddler and ingested could cause a serious reaction. Water bottles, infant formula, dry cereal (i.e. Cheerios) are acceptable.